

Isolation and structure elucidation of cyclopeptide alkaloids from the leaves of *Heisteria parvifolia*

Michel Boni Bitchi^{a,b}, Abdulmagid Alabdul Magid^b, Faustin Aka Kabran^a,
Philomène Akoua Yao-Kouassi^a, Dominique Harakat^c, Hamid Morjani^d, Félix Zanahi Tonzibo^{a,*},
Laurence Voutquenne-Nazabadioko^b

^a Laboratoire de Chimie Organique Biologique, UFR Sciences des Structures de La Matière et Technologie, Université Félix Houphouët-Boigny, 22 BP 582 Abidjan 22, Côte d'Ivoire

^b ICMR-UMR CNRS 7312, Equipe Chimie des Substances Naturelles, Campus Sciences, Bât. 18, BP 1039, 51687, Reims, France

^c Service Commun D'Analyses, Institut de Chimie Moléculaire de Reims (ICMR), CNRS UMR 7312, Bat. 18 B.P. 1039, 51687, Reims Cedex 2, France

^d BioSpecT EA7506, URCA, Faculté de Pharmacie, SFR CAP Santé, 1, Rue Du Maréchal-Juin, 51096, Reims, France

ARTICLE INFO

Keywords:

Heisteria parvifolia

Olacaceae

Cyclopeptide alkaloids

Cytotoxic activity

Chronic myeloid leukemia (K565) cell line

ABSTRACT

Heisteria parvifolia Sm. is prescribed in traditional medicine against numerous diseases in Côte d'Ivoire. Due to the shortcoming in scientific knowledge of use of this species, our investigations revealed five undescribed cyclopeptide alkaloids added to one known derivative namely anordianine. These compounds were elucidated by 1D and 2D-NMR experiments and comparison with literature data, and confirmed by HR-ESI-MS. Cytotoxic activity evaluation of these compounds against the chronic myeloid leukemia (K565) cell line exhibited an antiproliferative activity with cell growth inhibition from 13% to 46%.

1. Introduction

Cyclopeptide alkaloids are widespread and occur in several families: Asteraceae, Celastraceae, Euphorbiaceae, Fabaceae, Menispermaceae, Olacaceae, Pandaceae, Rhamnaceae, Rubiaceae, Sterculiaceae, and Urticaceae (El-Seedi et al., 2007; Gournelis et al., 1997; Morel et al., 2009; Tan and Zhou, 2006). Previous studies have reported cyclopeptides alkaloids from *Heisteria nitida* (El-Seedi et al. 1999, 2005). The cyclopeptide alkaloids *sensu stricto* were classified according to the number of amino acid constituents outside and the size of the macrocycle (inside) as 4(13); 5(13); 4(14) and 4(15) type of alkaloids (Joullie and Richard, 2004; Tan and Zhou, 2006). Several activities of cyclopeptides alkaloids have been reported, such as antimicrobial (Gournelis et al., 1997; Morel et al., 2005), insecticidal (Sugawara et al., 1996), sedative (Suh et al., 1997), and antiplasmodial activity (Suksamrarn et al., 2005).

The genus *Heisteria* belonging to the *Olacaceae* family comprises about 65 species in tropical America and 3 in Africa; namely *Heisteria parvifolia* Sm., *Heisteria trillesiana* Pierre ex Heckel, and *Heisteria zimmereri* Engl. *Heisteria parvifolia* Sm. is an evergreen shrub or small tree up to 15 (–20) m tall; 40 (–60) cm in diameter (Malaisse et al., 2004). *H. parvifolia* occurs from Senegal and south-western Mali eastward to

the Central African Republic and southward DR Congo and northern Angola; possibly also in Uganda and southern Sudan (Louppe et al., 2008). In Côte d'Ivoire, is locally abundant on sandy soils. Its wood is used for construction and tool handles. In several areas, the fruits are eaten fresh; the small oil-rich seeds are eaten fresh, roasted or cooked. The twigs are used as chew-sticks. Various *Heisteria* species are used by South-American Indians or in Africa in the treatment of rheumatism, abscesses, headache, throat infections, swellings, nose bleedings, pain in joints and muscles, diarrhea, hepatic infection (Kvist and Holm-Nielsen, 1987; Russo, 1992; Tan and Zhou, 2006). In traditional medicine in Ghana, ground roots of *H. parvifolia* are applied as enema against stomach-ache. In Congo, sap from the root bark is used as drops into the nose against migraine and into the eye to treat painful, infected eyes. Stem bark is taken in Ghana, in Côte d'Ivoire and DR Congo as cough medicine. In Gabon, bark is applied to circumcision wounds. In Ghana and Côte d'Ivoire, leaf decoctions are taken or applied as a bath to invigorate rachitic children and to treat convulsions. They are also used as analgesic and rubbed onto painful breasts of young mothers, and in Sierra Leone to treat tooth-ache. In Congo, leaf decoctions are administered against asthma, costal pain, stomach pain, and menstrual problems. Ground seeds are used to stupefy fish. In DR Congo, powdered bark is an ingredient in the preparation of arrow

* Corresponding author. Tel.: +225 05 07 19 67.

E-mail addresses: abdulmagid.alabdul-magid@univ-reims.fr (A.A. Magid), tonzibz@yahoo.fr (F.Z. Tonzibo).

poison (Abbiw, 1990; Burkill and families, 1997; Malaisse et al., 2004). Chemical investigations of *Heisteria* species have mainly revealed the presence of triterpenes and proanthocyanidines in *H. pallida* (Dirsch et al. 1992, 1993), cyclopeptide alkaloid in *H. nitida* (El-Seedi et al. 1999, 2005), scopolamine in *H. olivae* (Cairo-Valera et al., 1977), and acetylenic fatty acids in *H. accuminata* (Kraus et al., 1998). Up to date, only the composition of the seeds oil of *H. parvifolia* has been reported as mainly long-chain saturated fatty acids, oleic acid and other mono and di enoic fatty acids (Malaisse et al., 2004).

As a part of a continuing study for the discovery of medicinal Côte d'Ivoire species, five undescribed cyclopeptide alkaloids (1–5), together with one known compound (6), have been isolated and characterized from the leaves of *H. parvifolia*. Their cytotoxicity against the chronic myeloid leukemia K562 cells was evaluated.

2. Results and discussion

The crude alkaloid extract prepared with an acid-base method of air-dried and pulverized leaves of *H. parvifolia* was subjected to silica gel flash chromatography, eluted with increasingly polar mixtures of CHCl₃/MeOH. Further purification was performed using semi-preparative HPLC. As a result, five undescribed cyclopeptide alkaloids (1–5) were isolated and chemically characterized, together with one known cyclopeptide alkaloid, anordianine (6) (El-Seedi et al., 1999). Their structures (Fig. 1) were elucidated by 1D and 2D-NMR experiments and comparison with literature data, and confirmed by HR-ESI-MS.

The UV spectra of compounds 1–5 showed absorptions at 222–224 and 274–282 nm, wavelengths commonly assigned to peptide bonds and aromatic residues (Dongo et al., 1989; Kang et al., 2015; Schwing et al., 2011), while their IR spectra displayed bands at 3395 and 1682 cm⁻¹, which are typical of amide groups (Dongo et al., 1989; Schwing et al., 2011).




Fig. 1. Chemical structures of compounds 1–6.

Compound 1 was obtained as a white powder. Analysis of 1 by high-resolution electrospray ionization mass spectrometry (HR-ESI-MS) identified a pseudo-molecular ion [M + H]⁺ at *m/z* 457.2807, corresponding to the molecular formula C₂₅H₃₆N₄O₄ (calcd for C₂₅H₃₇N₄O₄, 457.2815), in combination with analysis of NMR data. The ¹³C NMR (Table 1) and HSQC spectra of 1 showed 25 carbon resonances for four methyls (δ_C 13.9, 16.7, 17.5, and 19.5), one *N*-methyl (δ_C 31.8), three methylenes (δ_C 23.3, 28.2, and 46.7), twelve methines (two of which were olefinic carbons at δ_C 116.7 and 124.9 and four were aromatic carbons *sp*² at δ_C 120.9, 121.9, 131.0, and 131.4), two quaternary aromatic carbons (δ_C 157.2 and 131.6), and three carbonyl carbons (δ_C 171.1, 167.5, and 165.2). The ¹H NMR spectrum (Table 1) displayed signals for two olefinic protons at δ_H 6.53 and 6.65, a singlet *N*-methyl (δ_H 2.68), four methyl doublets, four aromatic protons, and several methine and methylene protons. The NMR data of compound 1 (Table 1) showed great similarity with the NMR data previously reported for anordianine (compound 6) possessing a 14-membered ring type comprising a *p*-oxygenated *z*-styrylamine group (Dongo et al., 1989; El-Seedi et al. 1999, 2005). The presence of the *p*-oxygenated *z*-styrylamine group was indicated by two doublets at δ_H 6.53 and 6.65 (each 1H, *J* = 7.7 Hz; H-1 and H-2, respectively) corresponding to the *Z*-double bond and four aromatic protons appearing as doublets of doublets with *J*-values typical for an *o,m*-coupling pattern (H-12, 13, 15, and 16). The protons H-13 and H-15 (δ_H 7.11, dd, *J* = 8.7, 2.4 Hz) showed correlation with C-1 (δ_C 116.7) and H-12 and H-16 (δ_H 7.27, dd, *J* = 8.7, 2.4 Hz) with C-14 (δ_C 131.6) in HMBC spectrum. In cyclopeptide alkaloids, the H-9 (β-H of the β-hydroxy-amino acid moiety) chemical shift value (between 5.00 and 5.50 ppm) is characteristic. In this case, a doublets of doublets was present at δ_H 4.92 (*J* = 8.3, 1.5 Hz). In the COSY spectrum, two cross peaks were observed for H-9 (δ_H 4.92), more specifically with H-8 (δ_H 5.01, d, *J* = 8.3 Hz) and H-20 (δ_H 2.13, sept, *J* = 6.9 Hz). The proton signal of the CH-group in position 20 also showed cross peaks to H₃-21 (δ_H 1.32, d, *J* = 6.7 Hz) and H₃-22 (δ_H 1.05, d, *J* = 6.7 Hz). In the HMBC spectrum, correlations between C-9 (δ_C 83.4) and H-21 and H-22 were observed (Fig. 2). These data agreed with previously reported data for β-hydroxyleucine (anordianine). The methine and the methylene protons of proline were observed in the ¹H NMR spectrum: H-5 (δ_H 4.16, dd, *J* = 7.5, 1.9 Hz), H₂-17 (δ_H 1.65, m; 2.21, dd, *J* = 11.5–4.5 Hz), H₂-18 (δ_H 1.75, m; 1.95, m), and H₂-19 (δ_H 3.55, brt, *J* = 9.8 Hz; 3.85, m). In the COSY spectrum, cross peaks were observed between H-5/H-17, H-17/H-18, and H-18/H-19. In the HMBC spectrum, the H-5 exhibited correlations with C-19 (δ_C 46.7) and with the carbonyl C-4 (δ_C 167.5) (Fig. 2). The methyl doublets of *N*-methyl-valine (H-2' and H-3') appeared at δ_H 0.95 (*J* = 6.7 Hz) and 0.96 (*J* = 6.7 Hz) and the methine proton (H-1') of this moiety appeared as septuplet at δ_H 2.13 (*J* = 6.9 Hz). The HMBC experiment showed correlations between C-24 (δ_C 165.2) and H-25 (δ_H 3.57, d, *J* = 5.4 Hz) and H-1', and between the *N*-methyl carbon (δ_C 31.8) and the H-25 for this moiety. The combined use of 1D (¹H and ¹³C NMR) and 2D (COSY, HSQC, and HMBC) spectra allowed an unambiguous assignment of all protons and carbons of the amino acids units (β-hydroxyleucine, proline, *N*-methyl-valine residues) and the *p*-oxygenated *z*-styrylamine group (Table 1). Moreover, the connectivity between the constitutive parts of the molecule were ascertained by the HMBC correlations between the carbonyl C-4/H-2, the carbonyl C-7/H-5, the carbonyl C-24/H-8, and C-11/H-9 (Fig. 2). Moreover, the NOE relationships H-8/H-25, H-8/H-12 and H-9/H-16 agreed with the β-hydroxyleucine connection with the *N*-methyl-valine and the *p*-oxygenated *z*-styrylamine moieties whereas the NOE effects between H-8/H-19 agreed with the connection of β-hydroxyleucine with proline (Fig. 2). Compound 1 was named cycloheisterin A after its plant origin (Fig. 1).

Compound 2 displayed an [M + H]⁺ ion peak at *m/z* 471.2979 in the positive HR-ESI-MS, corresponding to the molecular formula C₂₆H₃₈N₄O₄, suggesting an additional methyl group compared to 1. The NMR spectroscopic data of 2 were almost identical with those of 1

Table 1
 ^{13}C NMR spectroscopic data for compounds 1–5 (500 MHz, CD_3OD).

	1		2		3		4		5	
	δ_{H} m (J in Hz)	δ_{C}	δ_{H} m (J in Hz)	δ_{C}	δ_{H} m (J in Hz)	δ_{C}	δ_{H} m (J in Hz)	δ_{C}	δ_{H} m (J in Hz)	δ_{C}
1	6.53, d (7.7)	116.7	6.53, d (7.7)	116.8	6.53, d (7.7)	116.3	6.53, d (7.7)	116.8	6.50, d (7.7)	116.8
2	6.65, d (7.7)	124.9	6.65, d (7.7)	125.0	6.66, d (7.7)	124.9	6.65, d (7.7)	124.9	6.60, d (7.7)	125.0
4	–	167.5	–	167.5	–	167.5	–	167.4	–	167.9
5	4.16, dd (7.5, 1.9)	62.5	4.14, d (7.5)	62.6	4.15, d (7.8)	62.5	4.12, d (8.1)	62.6	3.72, d (8.0)	62.4
7	–	171.1	–	171.0	–	171.1	–	171.0	–	170.7
8	5.01, d (8.3)	53.1	5.01, d (8.5)	52.0	5.01, d (8.3)	53.1	5.01, d (8.5)	53.0	4.82, (overlapped)	53.0
9	4.92, dd (8.3, 1.5)	83.4	4.93, dd (8.5, 1.9)	83.2	4.91, dd (8.3, 1.2)	83.4	4.92, dd (8.3, 1.3)	83.2	4.82, (overlapped)	82.7
11	–	157.2	–	157.2	–	157.2	–	157.2	–	157.1
12	7.27, dd (8.7, 2.4)	120.9	7.27, d (8.5)	120.8	7.27, d (8.1)	120.9	7.27, d (8.9)	120.8	7.20, m	119.6
13	7.11, dd (8.7, 2.4)	131.4	7.11, m	131.4	7.11, m	131.4	7.11, m	131.4	7.11, dd (8.5, 1.5)	131.3
14	–	131.6	–	131.8	–	131.6	–	131.6	–	131.3
15	7.11, dd (8.7, 2.4)	131.0	7.11, m	130.3	7.11, m	130.5	7.11, m	130.3	7.12, dd (8.5, 1.5)	130.2
16	7.27, dd (8.7, 2.4)	120.9	7.27, d (8.5)	120.8	7.27, d (8.1)	120.9	7.27, d (8.9)	120.8	7.20, m	120.3
17	1.65, m	28.2	1.67, m	28.2	1.65, m	28.1	1.64, m	28.2	1.54, m	28.6
	2.21, dd (11.5, 4.5)		2.21, dd (12.3, 4.9)		2.21, dd (12.1, 5.8)		2.22, dd (12.5, 6.1)		2.01, dd (11.3, 4.8)	
18	1.75, m	23.3	1.75, m	23.3	1.75, m	23.3	1.75, m	23.3	1.67, m	23.5
	1.95, m		1.97, m		1.95, m		1.97, m		1.82, m	
19	3.55, brt (9.8)	46.7	3.55, brt (10.8)	46.7	3.55, brt (9.8)	46.7	3.58, brt (9.3)	46.8	3.42, brt (9.2)	47.0
	3.85, m		3.90, ddd (10.8, 7.1, 3.2)		3.85, m		3.85, ddd (10.1, 9.3, 7.4)		3.71, m	
20	2.13, m	28.7	2.12, dq (6.8, 1.7)	28.7	2.13, m	28.7	2.11, m	28.7	2.10, m	28.6
21	1.32, d (6.7)	19.5	1.32, d (6.8)	19.4	1.32, d (6.8)	19.5	1.32, d (6.8)	19.4	1.27, d (6.8)	19.3
22	1.05, d (6.7)	13.9	1.07, d (6.8)	14.0	1.06, d (6.8)	13.9	1.07, d (6.8)	14.0	1.02, d (6.8)	14.0
24	–	165.2	–	164.3	–	165.3	–	164.4	–	165.0
25	3.57, d (5.4)	66.7	3.59, d (5.3)	72.7	3.62, d (5.0)	66.2	3.62, d (4.8)	72.2	4.10, dd (10.4, 4.5)	68.1
R ₁	N-CH ₃		N-CH ₃		N-CH ₃		N-CH ₃		N-CH ₃	
	2.68, s	31.8	2.92, s	40.3	2.66, s	31.8	2.91, s	39.7	2.97, s	41.1
R ₂			N-CH ₃				N-CH ₃		N-CH ₃	
			2.92, s	41.9			2.91, s	42.2	2.97, s	41.1
R ₃	Val		Val		iLeu		iLeu		Phe	
1'	2.13, m	30.2	2.45, m	27.2	1.88, m	36.8	2.16, m	33.9	3.12, dd (13.8, 10.6) 3.40, m	34.3
2'	0.95, d (6.7)	16.7	0.93, d (6.7)	15.1	1.02, m 1.47, m	25.1	0.77, m 1.42, m	26.3		134.1
3'	0.96, d (6.7)	17.5	0.97, d (6.7)	18.7	0.92, t (6.8)	10.3	0.93, t (6.9)	10.4	7.20, m	129.0
4'					0.96, d (6.8)	13.1	0.98, d (6.9)	11.5	7.30, m	128.5
5'									7.21, m	127.5
6'									7.30, m	128.5
7'									7.20, m	129.0

except for one additional methyl group (Table 1). The detailed analysis of the 2D-NMR spectra led to the identification, as in **1**, of the amino acids units (β -hydroxyisoleucine, proline, and valine residues) and the *p*-oxygenated *z*-styrylamine group (Table 1). The HMBC cross-peaks of the methyl signals at δ_{H} 2.92 (6H, s) to C-25 (δ_{C} 72.7) of the valine residue indicated that the terminal amino acid in **2** is *N,N*-dimethylvaline. Compound **2** was named cycloheisterin B.

Compound **3** displayed an $[\text{M} + \text{Na}]^+$ ion peak at m/z 493.2785 in the positive HR-ESI-MS, corresponding to the molecular formula $\text{C}_{26}\text{H}_{38}\text{N}_4\text{O}_4$, suggesting an additional methylene group compared to **1**. The ^1H and ^{13}C NMR values of **3** were almost superimposable on those of **1** (Table 1) excepting those of the *N*-methyl-valine residue in **1**. Instead, an *N*-methyl-isoleucine residue was identified as summarized in Table 1 (Tuenter et al., 2017). ^1H - ^1H COSY analysis confirmed the presence of an isoleucine residue. The HMBC cross-peak of H-8 of hydroxyisoleucine (δ_{H} 5.01) to C-24 of isoleucine residue (δ_{C} 165.3) and H-25 (δ_{H} 3.62) to C-24 and the *N*-methyl carbon (δ_{C} 31.8) to H-25 for this moiety confirmed that the terminal amino acid is *N*-methyl-isoleucine. Compound **3** was named cycloheisterin C (Fig. 1).

Cycloheisterin D (**4**) displayed an $[\text{M} + \text{H}]^+$ ion peak at m/z 485.3138 in the positive HR-ESI-MS, corresponding to the molecular formula $\text{C}_{27}\text{H}_{40}\text{N}_4\text{O}_4$, suggesting an additional methyl group compared to **3**. Comparing the NMR data of **4** with those of **3** (Table 1) and the analysis of the 2D-NMR spectra led to the identification, as in **3**, of the amino acids units (β -hydroxyisoleucine, proline and isoleucine residues) and the *p*-oxygenated *z*-styrylamine group (Table 1). The HMBC cross-peaks of the methyl signals at δ_{H} 2.91 (6H, s) to C-25 (δ_{C} 72.2) of the

isoleucine residue suggested that the terminal amino acid in **4** is *N,N*-dimethyl-isoleucine. Compound **4** was named cycloheisterin D (Fig. 1).

Cycloheisterin E (**5**) exhibited an $[\text{M} + \text{Na}]^+$ ion peak at m/z 541.2799 in the HR-ESI-MS spectrum, consistent with the molecular formula of $\text{C}_{30}\text{H}_{38}\text{N}_4\text{O}_4$. Comparing the NMR data of **5** with those of **1–4** (Table 1) and detailed analysis of the 2D-NMR spectra showed that it had the same macrocycle (inside) composed of the amino acids units (β -hydroxyisoleucine and proline) and the *p*-oxygenated *z*-styrylamine group (Table 1). The ^1H and ^{13}C -NMR spectra of **5** exhibited signals corresponding to an aromatic amino acid [δ_{H} 7.20–7.30, 5H]. Extensive 2D-NMR analysis enabled the full assignments of the *N,N*-dimethyl phenylalanine. (Tuenter et al., 2017). The presence of the *N,N*-dimethyl groups was confirmed by the HMBC correlation between the methyl signals at δ_{H} 2.97 (6H, s) and C-25 (δ_{C} 68.1) of the phenylalanine residue. The HMBC correlation between H-8 (δ_{H} 4.82) of the β -hydroxyisoleucine and the C-24 (δ_{C} 165.0) of the *N,N*-dimethyl phenylalanine confirmed it to be the terminal amino acid moiety. Compound **5** was named cycloheisterin E (Fig. 1).

The stereochemistry of the cyclopeptide alkaloids **1–6**, was proposed from the ^1H NMR coupling constants, ^{13}C NMR data, and NOESY analysis and by determining the absolute configuration of the amino acids by chiral HPLC after acid hydrolysis. With this purpose, compounds **1–6** were hydrolyzed and their amino acids analyzed through the chiral HPLC. In cycloheisterin A-E and **6**, proline has the *L* configuration and *N*-methyl-valine, *N,N*-dimethyl-valine, *N*-methyl-isoleucine, *N,N*-dimethyl-isoleucine, and *N,N*-dimethyl phenylalanine in cycloheisterin A-E, respectively and *N,N*-dimethyl-leucine in **6** were in




Fig. 2. Selected key HMBC and NOESY interactions for compound 1.

the *L* form. The ^{13}C -NMR chemical shift values of the α -amino acid of the macrocycle (proline in all five alkaloids) and the terminal units (*N*-methyl-valine in 1, *N,N*-dimethyl-valine in 2, *N*-methyl-isoleucine in 3, *N,N*-dimethyl-isoleucine in 4, *N,N*-dimethyl-phenylalanine in 5, and *N,N*-dimethyl-leucine in 6) match well with those previously reported for similar compounds and was in agreement with the fact that the majority of plant cyclopeptides are composed of *L*-amino acids (El-Seedi et al., 2005; Kang et al., 2015; Maldaner et al., 2011; Medina et al., 2016; Suksamrarn et al., 2005; Tuenter et al., 2017).

The configuration of the β -hydroxyleucine was established based on the available NMR data. In the case of the *erythro* form, $J_{\alpha,\beta}$ ca. 8.0 Hz, whereas for *threo* compounds $J_{\alpha,\beta}$ ca. 2.0 Hz (Fig. 3) (Dias et al., 2007; Gournelis et al., 1997; Mostardeiro et al., 2013; Tuenter et al., 2016). The coupling constant of the doublet corresponding to H-9 ($J_{\alpha,\beta}$) of compounds 1–5 ca. 8.3 Hz, clearly indicative of an *erythro* configuration. ^{13}C NMR spectroscopy is used for the elucidation of the absolute configuration of the β -hydroxy amino acids. For both *L-threo* and *D-threo* series, the signal of the α carbon appears at ca. δ_{C} 55.0, whereas for the β carbon, its signal appears at ca. δ_{C} 82.0 for the *D-threo* and ca. δ_{C} 86.0 for the *L-threo* (Fig. 3) (Mostardeiro et al., 2013). For the *L-erythro* series, the signal of the α carbon (C-8) appears at ca. δ_{C} 55.0, whereas for the *D-erythro* it appears at ca. δ_{C} 53.0. Important information is also observed for the β carbon (C-9): in the *L-erythro* series, the signal appears at ca. δ_{C} 81.5, whereas for the *D-erythro* configuration it appears at ca. δ_{C} 87.0 (Abu-Zarga et al., 1995; Caro et al., 2012; Dongo et al., 1989; Gournelis et al., 1997; Medina et al., 2016; Mostardeiro et al., 2013; Tuenter et al., 2016). These data show that the chemical shift of the β carbon is most indicative for the *L* and *D* forms of a β -hydroxy amino acids (Δ_{δ} 4–5 ppm) than α carbon (Δ_{δ} 0–3 ppm). The chemical shift of C-9 in compounds 1–5 was around $\delta_{\text{C-9}}$ 83.3, clearly suggestive for the *L-erythro* form, whereas the chemical shift of C-8 was around δ_{C} 53.0.




Fig. 3. Representatives and approximate NMR data for *threo* and *erythro* β -hydroxyleucine in cyclopeptide alkaloids.

Furthermore, the J value of the ^1H NMR signal attributed to the methyl group at position C-22 was 6.7 Hz, indicative for a pseudoaxial/equatorial coupling, typical for *L-erythro*- β -hydroxyleucine (Abu-Zarga et al., 1995; Gournelis et al., 1997; Tuenter et al., 2016). In addition, the cross-peak observed in the NOESY spectra of 1–5 between H-9 and H-20, H-9/H-21 and H-8/H-22 and the lack of the NOESY interaction between H-9 and H-8, suggests the *L-erythro* configuration for the β -hydroxyleucine moiety (Fig. 2). Furthermore, the NOESY effect observed between H-25 and H-1' indicated that these protons are co-facially oriented.

3. Conclusion

In summary, six compounds were isolated from the crude alkaloid extract of *H. parvifolia* leaves, among them five previously undescribed cyclopeptide alkaloids from the 4(14) type, 4 amino acid constituents outside and the 14-atoms of the macrocycle (inside). Their structures were established by different spectroscopic methods including 1D and 2D-NMR experiments as well as HR-ESI-MS analysis. Compound 6 (anoridianine) that has a unique substructure containing proline, was previously isolated from *Heisteria nitida* (El-Seedi et al., 1999). Compounds 1–5 were derivatives of anoridianine and differed in only the terminal amino acid which was *N*-methyl-valine in 1, *N,N*-dimethyl-valine in 2, *N*-methyl-isoleucine in 3, *N,N*-dimethyl-isoleucine in 4, and *N,N*-dimethyl-phenylalanine in 5. Cyclopeptide alkaloids have only been reported from a few families of the plant kingdom, in fact, they seem to be quite rare and present in small quantities. This kind of cyclopeptide alkaloids was isolated only in *Canthium anoridianum* (Rubiaceae) and *Heisteria nitida* (Olacaceae). Further phytochemical investigation on *Heisteria* species are needed to verify whether anoridianine derivative cyclopeptide alkaloids could be considered as a taxonomic markers for the genus *Heisteria*. The cytotoxic activity of compounds 1–6 against the chronic myeloid leukemia (K562) cell line was evaluated. Only compounds 2, 4 and 6 exhibited an anti-proliferative activity at the concentration 100 μM with cell growth inhibition of 46%, 44%, and 43%, respectively, whereas compounds 1, 3, and 5 showed cell growth inhibition of 13%, 19%, and 36%, respectively at the same concentration.

4. Experimental

4.1. General experimental procedures

Optical rotations were measured on a PerkinElmer model 341 polarimeter (589 nm, 20 °C). IR spectra were obtained on a Nicolet Avatar 320 FT-IR spectrometer with KBr disks. NMR spectra were acquired in CD₃OD on Bruker Avance DRX III 500 instruments (¹H at 500 MHz and ¹³C at 125 MHz). Standard pulse sequences and parameters were used to obtain 1D- (¹H and ¹³C) and 2D- (COSY, ROESY, HSQC and HMBC) NMR spectra. HR-ESI-MS experiments were performed using a Micromass Q-TOF high-resolution mass spectrometer (Manchester, UK). Mass spectra were recorded in the positive-ion mode in the range *m/z* 100–2000, with a mass resolution of 20000 and an acceleration voltage of 0.7 kV. Flash chromatography was conducted on a Grace Reveleris system equipped with dual UV and ELSD detection using Grace® cartridges (Silica gel or RP-18). A prepacked RP-C₁₈ column (Phenomenex 250 × 15 mm, Luna 5 μ) was used for semi-preparative HPLC. The eluting mobile phase consisted of H₂O with TFA (0.0025%) and CH₃CN with a flow rate of 5 mL/min and the chromatogram was monitored at 210, 250, 270, and 300 nm. TLC was performed on precoated silica gel 60 F₂₅₄ Merck and compounds were visualized by spraying the dried plates with Dragendorff's reagent.

4.2. Plant material

The leaves of *Heisteria parvifolia* Sm. were collected in Agboville forest in August 2016. They are identified by Pr. Akke Assi in the national center florestic of Félix Houphouët-Boigny University of Côte d'Ivoire (Ake assi 11049).

4.3. Extraction and isolation

The dried powdered leaves of *H. parvifolia* (1 kg) were wetted with 50% aq. NH₄OH (500 mL), macerated overnight and then percolated with 15 L of EtOAc. The organic solvent was concentrated under reduced pressure. The crude extract (26 g) was suspended in 2 L of EtOAc and extracted with an aqueous 2% H₂SO₄ solution (3 × 2 L). The acid phase was made alkaline with aqueous NH₃ and extracted with 3 × 2 L of CHCl₃. The CHCl₃ solution was washed with H₂O (2 L), dried (Na₂SO₄) and evaporated *in vacuo* to give 500 mg of crude alkaloid extract (yield 0.05%). The crude alkaloid extract was subjected to silica gel flash chromatography eluted with increasingly polar CHCl₃/MeOH (100:00–95:05) for 25 min, to yield 26 fractions (F1–26). Fractions F6, F8, F10, F12, F14 and F17 were subjected separately to semipreparative HPLC RP-18 chromatography, by eluting with an isocratic gradient (28% CH₃CN). Compound 4 (*t_R* 13.2 min, 31 mg) was obtained from fractions F6 and F8, compound 5 (*t_R* 14.9 min, 4 mg) from fraction F10, compound 6 (*t_R* 10.6 min, 6 mg) from fraction F12, compounds 2 (*t_R* 14.6 min, 6 mg) and 3 (*t_R* 17.3 min, 4 mg) from fraction F14, and compound 1 (*t_R* 11.3 min, 5 mg) from fraction F17.

4.3.1. Cycloheisterin A (1)

White amorphous powder; $[\alpha]_D^{20} = -148$ (c 0.5; MeOH); UV (MeOH) λ_{\max} (abs.) 222 (1.66), 274 (0.33); IR ν_{\max} 3395, 2972, 1682, 1508, 1205, 1133, 984, 720; ¹H and ¹³C NMR, see Table 1; HR-ESI-MS (positive ion mode) *m/z* 457.2807 [M + H]⁺ (calcd for C₂₅H₃₇N₄O₄, 457.2815).

4.3.2. Cycloheisterin B (2)

White amorphous powder; $[\alpha]_D^{20} = -187$ (c 0.52; MeOH); UV (MeOH) λ_{\max} (abs.) 222 (0.10), 282 (0.01); IR ν_{\max} 3439, 2969, 1681, 1508, 1204, 1136, 700; ¹H and ¹³C NMR, see Table 1; HR-ESI-MS (positive ion mode) *m/z* 471.2979 [M + H]⁺ (calcd for C₂₆H₃₉N₄O₄, 471.2971).

4.3.3. Cycloheisterin C (3)

White amorphous powder; $[\alpha]_D^{20} = -135$ (c 0.31; MeOH); UV (MeOH) λ_{\max} (abs.) 224 (1.38), 276 (0.37); IR ν_{\max} 3388, 2965, 1686, 1506, 1206, 1133, 985, 719; ¹H and ¹³C NMR, see Table 1; HR-ESI-MS (positive ion mode) *m/z* 493.2785 [M + Na]⁺ (calcd for C₂₆H₃₈N₄O₄Na, 493.2791).

4.3.4. Cycloheisterin D (4)

White amorphous powder; $[\alpha]_D^{20} = -179$ (c 0.23; MeOH); UV (MeOH) λ_{\max} (abs.) 222 (3.21), 280 (0.3); IR ν_{\max} 3395, 2972, 1682, 1508, 1205, 1133, 720; ¹H and ¹³C NMR, see Table 1; HR-ESI-MS (positive ion mode) *m/z* 485.3138 [M + H]⁺ (calcd for C₂₇H₄₁N₄O₄, 485.3128).

4.3.5. Cycloheisterin E (5)

White amorphous powder; $[\alpha]_D^{20} = -91$ (c 0.41; MeOH); UV (MeOH) λ_{\max} (abs.) 222 (0.91), 274 (0.5); IR ν_{\max} 3439, 2969, 1681, 1508, 1204, 1136, 700; ¹H and ¹³C NMR, see Table 1; HR-ESI-MS (positive ion mode) *m/z* 541.2799 [M + Na]⁺ (calcd for C₃₀H₃₈N₄O₄Na, 541.2791).

4.4. General procedure for determination of amino acid configurations

The absolute configurations of amino acids were determined by chiral HPLC after acid hydrolysis according to literature (Mostardeiro et al., 2013; Silva et al., 1996; Wang et al., 2017). Briefly, each solution of 1–5 (0.5 mg) in 6 N HCl (0.4 mL) was heated at 110 °C for 24 h and then concentrated to dryness. The residue was dissolved in H₂O (200 μL) to obtain the test solution, 10 μL of which was injected into chiral HPLC system with a Chiralpak IC column (250 mm × 4.6 mm I.D., 5 μm) maintained at 35 °C and detected at 254 nm: Isopropanol/*n*-hexane (90:10, v/v) containing 0.1% TFA was used as the mobile phase at a flow rate of 0.8 mL/min.

5. Cytotoxicity bioassay by MTS

K562 cells (chronic myeloid leukemia) were trypsinized, harvested, and spread onto 96-well flat-bottom plates at a density of 1000 cells per well, and then incubated for 24 h in RPMI 1640 Medium supplemented with 10% fetal bovine serum and antibiotics. After culture, the cells were treated with compounds 1–6 for 72 h. The cell cultures were then analyzed using 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium inner salt (MTS) according to the manufacturer's instructions (Promega Corporation, Charbonnières, France). Camptothecin was used as positive control. MTS is bioreduced by cells into a colored formazan product. Absorbance was analyzed at a wavelength of 540 nm with a Multiskan Ex microplate absorbance reader (Thermo Scientific, Paris, France). Percentage of cell growth was calculated as 100% × (absorbance of the treated cells)/(absorbance of the negative control cells). Control cells were treated with complete culture medium containing 0.2% DMSO. The values represent averages of three independent experiments.

Acknowledgements

The authors are grateful to Conseil Regional Champagne Ardenne, Conseil General de la Marne, Ministry of Higher Education, Research and Innovation (France) (MESRI) and EU-programme FEDER to the PIANET CPER project for financial support as well as the Ministry of Research of Côte d'Ivoire.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.phytochem.2019.112081>.

References

- Abbiw, D., 1990. Useful Plants of Ghana: West African Uses of Wild and Cultivated Plants. London. Intermediate Technology Publications, Royal Botanic Gardens, Kew.
- Abu-Zarga, M., Sabri, M., Al-Aboudi, A., 1995. New cyclopeptide alkaloids from *Ziziphus lotus*. J. Nat. Prod. 58, 504–511.
- Burkill, H.M., 1997. 2. The Useful Plants of West Tropical Africa, vol. 4. Kew Royal Botanic Gardens, pp. 284–285 families M-R.
- Cairo-Valera, G., De Budowski, J., Delle-Monache, F., Marini-Bettolo, G.B., 1977. A new psychoactive drug: *Heisteria olivae* (Olacaceae). Atti Accad. Naz. Lincei, Cl. Sci. Fis. Mat. Nat. Rend. 62, 363–364.
- Caro, M.S., de Oliveira, L.H., Ilha, V., Burrow, R.A., Dalcol, I.I., Morel, A.F., 2012. Absolute configuration of franganine. J. Nat. Prod. 75, 1220–1222.
- Dias, G.C., Gressler, V., Hoenzel, S.C., Silva, U.F., Dalcol, I.I., Morel, A.F., 2007. Constituents of the roots of *Melochia chamaedrys*. Phytochemistry 68, 668–672.
- Dirsch, V., Wiemann, W., Wagner, H., 1992. Anti-inflammatory activity of triterpene quinone-methides and proanthocyanidins from the stem bark of *Heisteria pallida*. Engl. Pharm. Pharmacol. Lett. 2, 184–186.
- Dirsch, V., Neszmélyi, A., Wagner, H., 1993. A trimeric propylargininid from stem bark of *Heisteria pallida*. Phytochemistry 34, 291–293.
- Dongo, E., Ayafor, J.F., Sondengam, B.L., Connolly, J.D., 1989. A new peptide alkaloid from *Canthium anordianum*. J. Nat. Prod. 52, 840–843.
- El-Seedi, H.R., Gohil, S., Perera, P., Torrsell, K.B.G., Bohlin, L., 1999. Cyclopeptide alkaloids from *Heisteria nitida*. Phytochemistry 52, 1739–1744.
- El-Seedi, H.R., Larson, S., Backlund, A., 2005. Chemosystematic value of cyclopeptide alkaloids from *Heisteria nitida* (Olacaceae). Biochem. Syst. Ecol. 33, 831–839.
- El-Seedi, H.R., Zahra, M.H., Göransson, U., Verpoorte, R., 2007. Cyclopeptide alkaloids. Phytochem. Rev. 6, 143–165.
- Gournelis, D.C., Laskaris, G.G., Verpoorte, R., 1997. Cyclopeptides alkaloids. Nat. Prod. Rep. 14, 75–82.
- Joullie, M.M., Richard, D.J., 2004. Cyclopeptide alkaloids: chemistry and biology. Chem. Commun. 18, 2011–2015.
- Kang, K.B., Ming, G., Kim, G.J., Ha, T.K., Choi, H., Oh, W.K., Sung, S.H., 2015. Jubanines F–J, cyclopeptide alkaloids from the roots of *Ziziphus jujuba*. Phytochemistry 119, 90–95.
- Kraus, C.M., Neszmélyi, A., Holly, S., Wiedemann, B., Nenninger, A., Torrsell, K.B., Bohlin, L., Wagner, H., 1998. New acetylenes isolated from the bark of *Heisteria acuminata*. J. Nat. Prod. 61, 422–427.
- Kvist, L.P., Holm-Nielsen, L.B., 1987. Ethnobotanical aspects of lowland Ecuador. Opera Bot. 92, 83–107.
- Louppe, D., Oteng-Amoako, A.A., Brink, M., 2008. In: Timbers, Plant Resources of Tropical Africa (Series), vol. 7. PROTA Foundation: Backhuys, Wageningen, pp. 385–386 1.
- Malaisse, F., N'Gasse, G., Lognay, G., 2004. *Heisteria parvifolia* (Olacaceae), an Underestimated shrub or small tree with Oil producing seeds. Syst. Geogr. Plants 74, 7–25.
- Maldaner, G., Marangon, P., Ilha, V., Caro, M.S.P., Burrow, R.A., Dalcol, I.I., Morel, A.F., 2011. Cyclopeptide alkaloids from *Scutia buxifolia* reiss. Phytochemistry 72, 804–809.
- Medina, R.P., Schuquel, I.T.A., Pomini, A.M., Silva, C.C., Oliveira, C.M.A., Kato, L., Nakamura, C.V., Santin, S.M.O., 2016. Ixorine, a new cyclopeptide alkaloid from the branches of *Ixora brevifolia*. J. Braz. Chem. Soc. 27, 753–758.
- Morel, A.F., Maldaner, G., Ilha, V., Missau, F., Silva, U.F., Dalcol, I.I., 2005. Cyclopeptide alkaloids from *Scutia buxifolia* Reiss and their antimicrobial activity. Phytochemistry 66, 2571–2576.
- Morel, A.F., Maldaner, G., Ilha, V., 2009. Cyclopeptide Alkaloids from higher plants. Alkaloids - Chem. Biol. 67, 79–141.
- Mostardeiro, M.A., Ilha, V., dahmer, J., Caro, M.S.B., Dalcol, I.I., Da Silva, U.F., Morel, A.F., 2013. Cyclopeptide alkaloids: stereochemistry and synthesis of the precursors of discarines C and D and myrianthine A. J. Nat. Prod. 76, 1343–1350.
- Russo, E.B., 1992. Headache treatments by native peoples of the Ecuadorian Amazon: a preliminary cross-disciplinary assessment. J. Ethnopharmacol. 36, 193–206.
- Schwing, K., Reyheller, C., Schaly, A., Kubik, S., Gerhards, M., 2011. Structural analysis of an isolated cyclic tetrapeptide and its monohydrate by combined IR/UV spectroscopy. ChemPhysChem 12 (10) 1981–1988.
- Silva, U.F., Cardoso, C.D., Zanatta, N., Icheln, D., Gehrcke, B., Morel, A.F., 1996. Determination of the stereochemistry of the *N,N*-dimethyl amino acid and the α -amino acid residue of peptide alkaloids by chiral gas chromatography. Phytochem. Anal. 7, 20–23.
- Sugawara, F., Ishimoto, M., Le Van, N., Koshino, H., Uzawa, J., Yoshida, S., Kitamura, K., 1996. Insecticidal peptide from mung bean: a resistant factor against infestation with azuki bean weevil. J. Agric. Food Chem. 44, 3360–3364.
- Suh, D.Y., Kim, Y.C., Kang, Y.H., Han, B.H., 1997. Metabolic cleavage of franguloline in rodents: *In vitro* and *in vivo* study. J. Nat. Prod. 60, 265–269.
- Suksamrarn, S., Suwannapoch, N., Aunchai, N., Kuno, M., Ratananukul, P., Haritakun, R., Jansakul, C., Ruchirawat, S., 2005. Ziziphine N, O, P and Q, new antiplasmodial cyclopeptide alkaloids from *Ziziphus oenoplia* var. *brunoniana*. Tetrahedron 61, 1175–1180.
- Tan, N.-H., Zhou, J., 2006. Plant cyclopeptides. Chem. Rev. 106, 840–895.
- Tuenter, E., Exarchou, V., Baldé, A., Cos, P., Maes, L., Apers, S., Pieters, L., 2016. Cyclopeptide alkaloids from *Hymenocardia acida*. J. Nat. Prod. 79, 1746–1751.
- Tuenter, E., Foubert, K., Staerk, D., Apers, S., Pieters, L., 2017. Isolation and structure elucidation of cyclopeptide alkaloids from *Ziziphus nummularia* and *Ziziphus spinachristi* by HPLC-DAD-MS and HPLC-PDA-(HRMS)-SPE-NMR. Phytochemistry 138, 163–169.
- Wang, X., Wu, H., Luo, R., Xia, D., Jiang, Z., Han, H., 2017. Separation and detection of free D- and L-amino acids in tea by off-line two-dimensional liquid chromatography. Anal. Method. 9, 6131–6138.